

*Mitteilungen aus den deutschen Schutzgebieten. Berlin. 20 Band.*

Ottweiler, Emil. Die Niederschlags-Verhältnisse von Deutsch-Südwestafrika. p. 1-84. [A compilation of all available data, including chart of mean annual rainfall and bibliography.]

*Naturwissenschaftliche Rundschau. Berlin. 22 Jahrgang. 21 März 1907.*

Süring, R. Wilhelm von Bezold. p. 153-155.

*Neueste Erdbeben-Nachrichten. Laibach. Jahrgang 6. 1906-7.*

Belar, A. Bodenbewegungen und die Stabilität der Bauten. p. 81-85.

Falsche Erdbebennachrichten. p. 103-105.

*Sitzungsberichte der Königlich preussischen Akademie der Wissenschaften. Berlin. 1907. XII.*

Warburg, E. and Leithäuser, G. Ueber die Oxydation des Stickstoffs bei der Wirkung der stillen Entladung auf atmosphärische Luft. p. 229-234.

*Weltall (Das). Berlin. 7 Jahrgang. 1907, April 1.*

Linke, Felix. Vom Staube. p. 193-201.

Krebs, Wilhelm. Witterungsvoraussicht und Sonnentätigkeit. p. 201-203.

*Wetter (Das). Berlin. 24 Jahrgang. März 1907.*

Großmann, —. Die Wettervorhersage auf den Witterungsberichten des Königl. sächsischen meteor. Institutes. p. 49-54.

Kaiser, Max. Historische Entwicklung unserer Kenntnis der Land- und Seewinde auf der Erde und Darstellung der gegenwärtigen Theorien. p. 54-65.

Wahlburg, Eduard Schiefer Edler von. Die Antizyklone der letzten Januar-Dekade 1907. p. 68-72.

*Atti della Reale accademia dei Lincei. Roma. v. 15. 1906.*

Monti, V. Sulla misura della velocità di propagazione delle perturbazioni sismiche in rapporto alla sismometria razionale. p. 15-18.

Pizzetti, —. Intorno al calcolo della rifrazione astronomica, senza speciali ipotesi sul modo di variare della temperatura dell'aria coll'altezza. p. 73-81.

Chella, Silvio. Misura del coefficiente di attrito interno dell'aria a basse temperature. p. 119-125.

Chistoni, Ciro. Risultati pireliometrici ottenuti dal 22 agosto a tutto giugno 1903 al R. Osservatorio geofisico di Modena. p. 126-132.

Monti, V. Sulla probabile origine della distribuzione dei temporali italiani a seconda delle stagioni. p. 173-175.

Chistoni, Ciro. Misure pireliometriche eseguite sul Monte Cimone nell'estate del 1902 e nell'estate del 1903. p. 208-213.

Teglio, Emilio. Contributo allo studio del pireliometro a compensazione elettrica dell'Angström. p. 214-216.

Monti, V. Sull'interpretazione matematica dei sismogrammi. p. 217-219.

Chistoni, Ciro. Misure pireliometriche eseguite sul Monte Cimone nell'estate del 1904 e nell'estate del 1905. p. 276-281.

Eredia, Filippo. La pioggia a Roma. p. 450-456. [Gives rainfall in extenso, by months, 1825-1906.]

Blaserna, —. Sulle esperienze degli spari contre la grandine, eseguite a Castelfranco Veneto negli anni 1902-1906. p. 680-682.

*Revista marittima. Roma. Anno 41. Marzo 1907.*

Penne, Renzo de la. Osservazioni di scariche elettriche dell'atmosfera. [Observations made aboard the *Calabria* in a voyage around the world.]

*Società geografica Italiana. Roma. Serie 4. Vol. 7.*

— La regione meno piovosa delle Alpi. (Feb., 1906.) p. 142-143.

— La rete meteorologica eritrea. (Feb., 1906.) p. 155.

Baratta, Mario. I terremoto di Calabria. (Maggio 1906.) p. 432-459.

Alessandri, Camillo. Due mesi sulla vetta del Monte Rosa. (Luglio 1906.) p. 639-662. [Account of Regina Margherita observatory, Monte Rosa.]

Tancredi, A. M. Nota sul clima del Seralhè (Colonia Eritrea). (Dic. 1906.) p. 1192-1249. [Includes collected data for Adi-Ugri and Chenafenah.]

Baratta, M. Sulla distribuzione topografica dei terremoti nel Chili. (Gen. 1907.) p. 30-37.

*Boletín de la Sociedad geográfica de Lima. Año 16. Tome 19.*

Victoria, Ernesto G. Evaporación y frío producido por ella en Lima. p. 1-58.

Castre, Emilio. El departamento de San Martín y nuestras regiones orientales. Conferencia dada en la Sociedad geográfica de Lima. p. 59-97. [Climate, p. 61-62.]

*Hemel en dampkring. Den Haag. 4. Jaargang.*

Kater, J. De groene straal bij op en ondergang der zon. p. 170-171.

M. Het fiebiel van hooge drukking in de tweed helft van Januari dezes jaars. p. 171-173. [Includes charts Jan. 19-24.]

M. Weerkundige waarnemingen te de Bilt 1897-1906. p. 173-176. [Brief résumé 1897-1906.]

## NORTH ATLANTIC WEATHER.

By MR. JAMES PAGE, Chief of the Division of Ocean Meteorology.

[Compiled from the daily observations, at Greenwich mean noon, furnished by cooperating observers at sea.]

The distribution of pressure over the North Atlantic Ocean at the instant of Greenwich mean noon of March 1 and the attendant circulation of the winds is shown on Chart X. Pressure is above the average over Iceland, 29.80 inches, and a marked anticyclonic area covers the eastern portion of the ocean and the western shores of the Continent of Europe. The latter feature of the pressure remained practically constant thruout the entire month, the minimum barometric reading at Ponta Delgada, Azores, during the period March 1-30 being 30.18 inches, recorded March 25, consequent upon a northward recession of the center of the high to the British Isles. Over the latter upon the same date readings of 30.40 inches and upward were recorded.

As a result of this special distribution of pressure the northeast trades blew thruout the entire month without interruption, extending in a continuous belt from the latitude of Cape Finisterre to the Line. At the entrance to the Channel and westward along the transatlantic routes as far as the meridian of 20° the winds blew almost continuously from the southwest quadrant, and fair weather prevailed. Upon one day only, viz, March 16, did these winds attain gale force, the result of a cyclonic storm central at the time in the vicinity of the Faroes, where the recorded pressure upon the date mentioned was 29.00 inches.

Over the western half of the ocean variations of pressure succeeded one another with marked rapidity, and as a consequence severe weather was the rule. The high central over New England yielded March 2 and was succeeded March 3 by a shallow low which extended eastward to mid-ocean. Upon this date the transatlantic routes between the meridians of 60° W. and 35° W. were accordingly visited by westerly gales, without, however, any decided change in the barometer.

Pressure over Iceland fell from 29.50 inches on March 2 to 28.90 inches on March 5, with the result that vessels following the route north about Scotland experienced southwesterly gales of force 10 and 11 during this period.

On March 5 a feeble area of low pressure moved eastward across Hatteras and on March 6 was central with a well developed system of cyclonic winds in the neighborhood of 37° N., 66° W., the lowest recorded pressure being 29.30 inches. From this point it moved northeastward, developing into a hurricane during the early hours of March 7.

The distribution of pressure and the circulation of the winds at Greenwich mean noon of this date are shown on Chart XI, the storm at this hour being central in latitude 42° N., longitude 56° W. Of the large number of vessels which experienced the hurricane's severity, the *Pretoria*, (German S. S., Schröter, report by 3d officer Suppeln), appears to have most nearly approached the center. The vessel was bound from Hamburg to New York, and at Greenwich mean noon of March 6 found herself in latitude 41° 55' N., longitude 53° 20' W. The remarks of the observer from this time forward are as follows:

"At 2 p. m. of March 6 the wind went from west, 2, to the southward, and so continued until 3 a. m. of March 7, with heavy rain squalls; lightning and thunder covering the whole sky. The wind then shifted to WSW., force 11, and at 6 a. m. to W., force 12, with a heavy hail squall. The lowest barometer, 725.0 millimeters (28.54 inches), occurred at 2 a. m., the position at the time being latitude 41° 51' N., longitude 57° 00' W., and the wind south, force 11."

The *Brandenburg*, (German S. S., Woltersdorff, Bremen to New York, report by officer Jaehnigen), also found herself within dangerous proximity to the storm center, altho in the

opposite semicircle. The position of the vessel at Greenwich mean noon of March 6 was latitude  $41^{\circ} 19' N.$ , longitude  $57^{\circ} 51' W.$ —about 180 miles, therefore, to the westward of the *Pretoria*. The shifts of the wind during the advance of the storm were ENE., 10; NNW., 11; WNW., 10. The observer reports: "Continuous violent rain until 2 p. m. of the 6th; a tremendous sea running, causing the ship to labor fearfully; the wind blows at times with hurricane force; at 6 p. m. the sea was so very boisterous that the ship had no steerage way; both engines were accordingly stopt; at 3 a. m. of March 7, the storm had abated sufficiently to allow us to resume our course; lowest barometer, 733 millimeters (28.88 inches), at midnight".

The center of the storm moved northward across Cape Race during March 8, and conditions were thence undisturbed until March 11. Upon this date another depression appeared in the neighborhood of  $39^{\circ} N.$ ,  $65^{\circ} W.$ , and, moving northeastward, was followed by northwesterly gales in the region to the southward of Nova Scotia. This was followed by a period of quiet weather which terminated on the 19th. Upon the last-named date a low moved eastward from the Great Lakes and on March 20 was central over the Bay of Fundy, the barometer at Eastport, Me., reading at 8 a. m. 28.66 inches. Southerly gales of force 10 and 11 swept the transatlantic routes from the American coast to the meridian of  $60^{\circ}$  west. The center of the depression did not at any time come within the region of observation at sea.

On March 24 an elongated trough of low pressure extended southeastward from Cape Race to a point situated in latitude  $35^{\circ} N.$ , longitude  $40^{\circ} W.$  On the western slope of this trough northwesterly gales of force 8 to 9 prevailed, covering a belt

300 miles in width. On the eastern slope southerly and southeasterly winds of force 6 were the rule, rising to force 8 thruout a limited area at the southern extremity. As the day advanced the axis of this trough assumed a more easterly direction, the trough itself at the same time increasing in depth, with the result that thruout March 25 the transatlantic routes from the longitude of Cape Race to the meridian of  $30^{\circ} W.$ , were the scene of steady southwesterly and westerly gales of force 9 and 10.

On the 26th of the month a tropical depression made its appearance between Bermuda and Porto Rico, in which the *Epsom* (British S. S., Cox, Channel to Galveston, report by officer Williams) and the *Tampico* (British S. S., Westcott, Channel to New Orleans, report by officer Haworth) became involved on the 27th. According to the report of the former vessel the slow initial fall of the barometer which marks the approach of storms of this nature set in at noon of March 25. At 4 a. m. of the 26th, the barometer rose slightly and the wind became variable, finally settling in the northeast, while a heavy northwesterly swell at the same time made itself felt. The position of the vessel at Greenwich mean noon was latitude  $33^{\circ} N.$ , longitude  $69^{\circ} W.$ ; wind NE., barometer 29.44 inches. Fifteen minutes later a squall of wind heralded the break of the threatening gale from the north. Fierce squalls of hurricane force were frequent and a very high and dangerous sea soon rose. The hurricane continued to rage thruout the day, the barometer meanwhile rising, altho very slowly. At Greenwich mean noon of the 27th the position of the *Epsom* was latitude  $30^{\circ} 30' N.$ , longitude  $63^{\circ} 10' W.$ , wind N., 12; barometer 29.71 inches, weather overcast and squally. At 1 p. m. the sky cleared and the wind and sea soon moderated.

## THE WEATHER OF THE MONTH.

By Mr. P. C. DAY, Assistant Chief, Division of Meteorological Records.

### PRESSURE.

The distribution of mean atmospheric pressure for March 1907, over the United States and Canada is graphically shown on Chart VI, and the average values and departures from the normal are shown for each station in Tables I and V.

During March, 1907, the distribution of the average pressure showed two well marked variations from the normal. Pressure averaged unusually low over all northwestern districts of the United States and Canada, and relatively high over all southern and eastern districts.

As a result of this reversal of pressure distribution, no high-pressure areas of any considerable magnitude moved southward from their usual source of origin over the Great Plains region east of the Canadian Rockies, and such cold waves as overspread the more northern districts were correspondingly lacking in severity.

On the other hand, the presence of unusually high pressure over the Gulf of California and adjacent territory and all southern and eastern districts forced the warm southerly winds of those latitudes far to the north of their usual limits.

Over the upper Missouri Valley and thence westward to the Pacific and northward over the Canadian Provinces the pressure was almost continuously below the average, while over southern California, Arizona, and New Mexico comparatively high pressure was the rule during most of the month.

Pressure also averaged considerably above the normal in all districts along the Gulf and Atlantic coasts. The entire region west of the Rocky Mountains, the Great Plains south of Nebraska, the lower Mississippi Valley, and the Gulf States were not within the direct influence of any considerable area of high pressure, having its origin in northern districts, during the month; on the other hand, no portion of the country was exempt from the influence of the numerous low areas which moved eastward from the Pacific.

### TEMPERATURE.

The warm waves that overspread nearly all districts of the United States east of the Rocky Mountains from the 18th to 23d and from the 24th to 29th established new records at many points for both the highest monthly mean and the highest maximum temperatures ever recorded in March at the respective stations.

The abnormally warm weather during the above periods occurred in connection with marked depressions of the barometer in the central districts, but generally without the usual cloud formations and attended by little or no precipitation, which, with comparatively high pressure over the Gulf States and in the extreme Southwest, gave warm southerly winds and almost midsummer temperatures over the Great Plains, central valleys, and all eastern districts. Maximum temperatures far in excess of any previous March record, and in some sections higher than before recorded in any previous April or May, were recorded about the 23d, and again about the 29th.

The month as a whole was one of marked temperature excess over all portions of the United States, except the extreme eastern portion of Maine and over the Pacific coast districts.

Over practically all that portion of the United States from the Rocky Mountains eastward the average for the month exceeded the normal by more than  $6^{\circ}$  daily, and over the greater portion of the central Mississippi Valley region, Kansas, Oklahoma, and Texas the normal was exceeded by more than  $10^{\circ}$  daily.

Over the greater portion of the territory between the Rocky Mountains and the Mississippi Valley and south of Nebraska, the temperature has averaged continuously above the normal for the four months from December to March, inclusive. The average excess during that period over portions of Kansas, Oklahoma, Texas, Louisiana, and Arkansas, ranges from  $7^{\circ}$  to more than  $10^{\circ}$  daily.